

AD-A144 716

QUASISTABLE PARAMETRIC OPTIMIZATION WITHOUT COMPACT
LEVEL SETS(U) WISCONSIN UNIV-MADISON MATHEMATICS
RESEARCH CENTER L MCLINDEN JUN 84 MRC-TSR-2708

1/1

UNCLASSIFIED


DAAG29-80-C-0041

F/G 12/1

NL



END
DATE
FILMED
9-84
DTIC



MICROCOPY RESOLUTION TEST CHART
 NATIONAL BUREAU OF STANDARDS-1963-A

AD-A144 716

MRC Technical Summary Report #2708

QUASISTABLE PARAMETRIC OPTIMIZATION
WITHOUT COMPACT LEVEL SETS

L. McLinden

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

June 1984

(Received November 22, 1983)

AUG 27 1984

A

**Approved for public release
Distribution unlimited**

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, D. C. 20550

84 08 24 039

DTIC FILE COPY

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

QUASISTABLE PARAMETRIC OPTIMIZATION WITHOUT COMPACT LEVEL SETS

L. McLinden

Technical Summary Report #2708
June 1984

ABSTRACT

The well-known perturbational duality theory for convex optimization is refined to handle directly, in locally convex Hausdorff spaces, problems involving noncoercive convex functionals together with unbounded densely defined linear operators or, more generally, convex processes. The theory presented includes conjugacy, recession, and ϵ -subdifferential formulas for the two fundamental pairs of dual operations and also includes systematic treatment of ϵ -solutions.

AMS (MOS) Subject Classifications: Primary: 49-00, 49A27, 49B27;
Secondary: 65K10, 90C25, 90D05

Key Words: Convex optimization, noncoercive functionals, unbounded operators, convex processes, ϵ -solutions, weak local compactness, recession theory, dual operations, ϵ -subdifferentials, saddlepoint problems, perturbational duality, abstract Hamiltonian system, abstract Euler-Lagrange system

Work Unit Number 5 (Optimization and Large Scale Systems)



Sponsored by the United States Army under Contract No. DAAG29-80-C-0041, and the National Science Foundation under Grant No. MCS-8102684.

A1

SIGNIFICANCE AND EXPLANATION

Dual variational principles, which arise classically in physics and mechanics and typically involve partial differential operators, usually correspond to dual pairs of constrained convex optimization problems in various infinite-dimensional function space settings. Constrained optimization denotes the search for an optimum, say a minimum, of some given criterion functional over a set of candidate solutions which satisfy given side conditions, or constraints. Convex problems form the simplest class of nonlinear problems. Dual pairs of convex optimization problems are nonlinear generalizations of the familiar, widely useful dual pairs of linear programming problems. Dual pairs of constrained convex optimization problems arise also in fields other than physics and mechanics, including mathematical economics, operations research, management science, many subfields of engineering, and mathematics itself.

The mathematical model for constrained convex optimization problems known as perturbational duality theory, developed some 10 to 15 years ago, provides a nearly complete mathematical treatment of those basic situations which involve "nice" (i.e. coercive) convex functionals and "nice" (i.e. bounded) linear operators. Much less developed is the treatment for problems involving noncoercive functionals and unbounded operators only densely defined. Even less understood are problems involving more complicated operators called convex processes, which in general are multivalued and not even densely defined, and which are important in problems from, e.g., mathematical economics.

This paper presents a mathematical theory, valid in very general spaces, for dual pairs of constrained convex optimization problems involving noncoercive functionals and unbounded linear operators or, more generally, convex processes. It constitutes a fundamental refinement of the well-known perturbational duality model cited above. Systematic treatment of approximate optimal solutions is also provided, rendering the model more amenable to further, numerical analysis.

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

QUASISTABLE PARAMETRIC OPTIMIZATION WITHOUT COMPACT LEVEL SETS

L. McLinden

It is well known that a lower semicontinuous (l.s.c.) function $\phi : X \rightarrow (-\infty, \infty]$, $\phi \not\equiv \infty$, achieves a finite infimum if the sets $\{x | \phi(x) \leq \alpha\}$, $\alpha \in \mathbb{R}$, are compact. Convex conjugate duality establishes that, if ϕ is convex, a dual sufficient condition for such compactness is that the conjugate function ϕ^* be continuous at 0. Much work in constrained optimization has dealt with elaborating and implementing this result (see References). The present work is rooted in the following generalization: a direct sufficient condition for such a convex ϕ to achieve a finite infimum is that (i) $\{x | \phi(x) \leq \alpha\}$, $\alpha \in \mathbb{R}$, are locally compact and (ii) ϕ is constant along the whole line generated by any direction in which ϕ does not eventually increase. Here we present theorems embodying this result in strengthened, augmented form tailored to basic structure occurring widely in constrained optimization. They go well beyond the closest counterparts now available (including, notably, [5], [10], [9], [1, Ch. 14]).

A word on notation. All spaces are locally convex Hausdorff topological vector spaces over \mathbb{R} ; X, V are paired in duality, as are U, Y . For a multifunction T from Z into W (i.e. $T : Z \rightarrow 2^W$), $G(T) := \{(z, w) | w \in T(z)\}$, $D(T) := \{z | T(z) \neq \emptyset\}$, $R(T) := \{w | \exists z, w \in T(z)\}$. For T and any $\phi : Z \rightarrow [-\infty, \infty]$, define $T\phi$ on W by $(T\phi)(w) := \inf\{\phi(z) | z \in T^{-1}(w)\}$, with $\inf \emptyset := \infty$. For ϕ l.s.c. proper convex, $\phi_0^+(z) := \sup\{\lambda^{-1}(\phi(\bar{z} + \lambda z) - \phi(\bar{z})) | 0 < \lambda < \infty\}$ for any fixed $\bar{z} \in \text{dom } \phi := \{z | \phi(z) < \infty\}$, and if Z is paired in duality with W , $\phi^*(w) := \sup\{\langle z, w \rangle - \phi(z) | z \in Z\}$ and $G(\partial_\epsilon \phi) := \{(z, w) | \phi^*(w) \leq \epsilon + \langle z, w \rangle - \phi(z)\}$ for each fixed $\epsilon \in [0, \infty)$. We abbreviate "weakly locally compact" by w.l.c.

THEOREM I. Let h be l.s.c. proper convex on X with $k := h^*$. Let A be a densely defined linear operator from X into U with closed graph and adjoint B . Define kB on Y by $(kB)(y) := k(By)$ if $y \in D(B)$ and $:= \infty$ if $y \notin D(B)$. Assume (i) $D(A) \cap \text{dom } h$ is nonempty and w.l.c. and (ii) $x \in A^{-1}(0)$, $h0^+(x) < 0 \implies h0^+(-x) < 0$. Then (1) $(kB)^*(u) = (Ah)(u) > -\infty$ and (2) $(Ah)0^+(u) = (A(h0^+))(u) > -\infty$, with both infima on the right attained whenever not vacuously. For any $\varepsilon \in [0, \infty)$, $y \in \partial_\varepsilon(Ah)(u) \iff u \in \partial_\varepsilon(kB)(y) \iff y \in D(B)$, $u \in A\partial_\varepsilon k(By)$.

Notice (1) $\implies \emptyset \neq R(B) \cap \text{dom } k$. Further, by (1), (2) is equivalent to:

$$(2') \quad \sup_{y \in B^{-1} \text{dom } k} \langle u, y \rangle = \inf_{x \in A^{-1}(u)} \sup_{v \in \text{dom } k} \langle x, v \rangle > -\infty,$$

with the infimum attained whenever $u \in R(A)$. If (ii) holds in the form

" $x \in A^{-1}(0)$, $h0^+(x) < 0 \implies x = 0$," the infimum problems on the right of (1), (2),

(2') have weakly compact ε -approximate solution sets for all $\varepsilon \in [0, \infty)$.

"Quasistable" in the title denotes the fact that, whenever $(Ah)(u) < \infty$ in (1), one can prove the u.s.c. proper concave $q(v) := \inf\{h(x) - \langle x, v \rangle \mid x \in A^{-1}(u)\}$ on V is Mackey quasicontinuous [5] and that the "inf" defining q is attained for all v (e.g., $v = 0$) in the Mackey relative interior of $\{v \mid q(v) > -\infty\}$. Counterparts of these remarks apply below.

THEOREM II. For $j = 1, \dots, m$ let f_j be l.s.c. proper convex on X with $g_j := f_j^*$. Assume (i) $\text{dom } f_2, \dots, \text{dom } f_m$ are w.l.c. and (ii) $x_1 + \dots + x_m = 0$, $f_1 0^+(x_1) + \dots + f_m 0^+(x_m) < 0 \implies f_1 0^+(-x_1) + \dots + f_m 0^+(-x_m) < 0$. Then (1) $(g_1 + \dots + g_m)^*(x) = (f_1 \square \dots \square f_m)(x) > -\infty$, and (2) $(f_1 \square \dots \square f_m)0^+(x) = (f_1 0^+ \square \dots \square f_m 0^+)(x) > -\infty$, with both infima on the right attained. [\square denotes inf-convolution.] For any $\varepsilon \in [0, \infty)$, $v \in \partial_\varepsilon(f_1 \square \dots \square f_m)(x) \iff x \in \partial_\varepsilon(g_1 + \dots + g_m)(v) \iff x \in \bigcup \{\partial_{\varepsilon_1} g_1(v) + \dots + \partial_{\varepsilon_m} g_m(v) \mid \varepsilon_j > 0, \varepsilon_1 + \dots + \varepsilon_m = \varepsilon\}$.

THEOREM III. Let h be l.s.c. proper convex on X with $k := h^*$. Let A be a closed convex process [8, §39] from X into U (i.e. $G(A)$ is a nonempty closed

convex cone) with adjoint A^* from Y into V given by $v \in A^*(y) \iff \langle u, y \rangle \leq \langle x, v \rangle$ for all $u \in A(x)$. Assume (i) $\emptyset \neq D(A) \cap \text{dom } h$ and $\text{dom } h$ is w.l.c. and (ii) $x \in A^{-1}(0)$, $h^+(x) \leq 0 \implies h^+(-x) \leq 0$. Then, for $B := A^{*-1}$, (1) $(Bk)^*(u) = (Ah)(u) > -\infty$ and (2) $(Ah)^0(u) = (A(h^0))(u) > -\infty$, with both infima on the right attained whenever not ∞ vacuously. Further, $y \in \partial_0(Ah)(u) \iff u \in \partial_0(Bk)(y) \iff [\exists x \in A^{-1}(u) \exists v \in B^{-1}(y) \text{ such that } \langle u, y \rangle > \langle x, v \rangle, v \in \partial_0 h(x)]$. For any $\epsilon, \delta \in [0, \infty)$ and $(v, y) \in G(B)$ with $k(v) < \delta + (Bk)(y) < \infty$, $y \in \partial_\epsilon(Ah)(u) \iff u \in \partial_\epsilon(Bk)(y) \implies [u \in R(A) \text{ and } \exists x \in A^{-1}(u) \exists \alpha > 0 \exists \beta > 0 \text{ such that } \alpha + \beta = \epsilon + \delta, \langle u, y \rangle > \langle x, v \rangle - \beta, x \in \partial_\alpha k(v)] \implies u \in \partial_{\epsilon+\delta}(Bk)(y) \iff y \in \partial_{\epsilon+\delta}(Ah)(u)$.

THEOREM IV. Let F be l.s.c. proper convex on $X \times U$. Let A be a linear transformation from X into U with either A continuous or A densely defined with closed graph. In the latter case, assume $[F(\cdot, u) \not\equiv \infty \implies F(\cdot, u) \text{ somewhere continuous}]$, or dually, $[F^*(v, \cdot) \not\equiv \infty \implies F^*(v, \cdot) \text{ somewhere continuous}]$. Assume (i) $\{x \in D(A) \mid \exists u, F(x, u) < \infty\}$ is nonempty and w.l.c. and (ii) $x \in D(A)$, $F^0(x, Ax) \leq 0 \implies F^0(-x, -Ax) \leq 0$. Then

$$-\infty < \min_{x \in D(A)} F(x, u + Ax) = \sup_{y \in D(A^*)} \{-F^*(A^*y, -y) - \langle u, y \rangle\} =: p(u),$$

$$-\infty < \min_{x \in D(A)} \sup_{(y, v) \in \Delta} \langle (x, u), (v, -y) \rangle = \sup_{y \in \Delta_0} \langle u, -y \rangle = p^0(u),$$

where $\Delta := \{(y, v) \mid y \in D(A^*), F^*(A^*y, -y) < \infty\}$, $\Delta_0 := \{y \mid (y, 0) \in \Delta\}$. For any $\epsilon \in [0, \infty)$, $-y \in \partial_\epsilon p(u) \iff y \in \text{Arg } \epsilon\text{-max}\{-F^*(A^*\hat{y}, -\hat{y}) - \langle u, \hat{y} \rangle \mid \hat{y} \in D(A^*)\} \iff [y \in D(A^*) \text{ and } \exists x \in D(A) \text{ such that } (A^*y, -y) \in \partial_\epsilon F(x, u + Ax)]$, and any such x belongs to $\text{Arg } \epsilon\text{-min}\{F(\hat{x}, u + A\hat{x}) \mid \hat{x} \in D(A)\}$. If also (iii) Δ_0 is w.l.c. then, for each u satisfying $[y \in D(A^*), F^0(A^*y, -y) + \langle u, y \rangle \leq 0 \implies F^0(-A^*y, y) - \langle u, y \rangle \leq 0]$, one has $p(u) \in \mathbb{R}$ and the "sup" defining $p(u)$ is actually "max."

THEOREM V. Let H be closed proper convex-concave [9] on $X \times Y$, put $F(x, u) := \sup\{H(x, y) - \langle u, y \rangle \mid y \in Y\}$, and assume this F together with A satisfy

the hypotheses of Theorem IV. Then

$$-\infty < \min_{x \in D(A)} \sup_y \{H(x,y) - \langle Ax, y \rangle - \langle u, y \rangle\}$$

$$= \sup_{y \in D(A^*)} \inf_x \{H(x,y) - \langle x, A^* y \rangle - \langle u, y \rangle\} =: p(u),$$

$$-\infty < \min_{x \in D(A)} \sup_{(y,v) \in \Delta} \langle (x,u), (v,-y) \rangle = \sup_{y \in \Delta_0} \langle u, -y \rangle = p_0^+(u),$$

where $\Delta := \{(y,v) | y \in D(A^*), G(y, v + A^* y) > -\infty\}$, $\Delta_0 := \{y | (y,0) \in \Delta\}$ for $G(y,v) := \inf\{H(x,y) - \langle x, v \rangle | x \in X\}$. For any $\varepsilon \in [0, \infty)$, $-y \in \partial_\varepsilon p(u) \iff$

$$y \in \text{Arg } \varepsilon\text{-max}_{\hat{y} \in D(A^*)} \inf_{\hat{x}} \{H(\hat{x}, \hat{y}) - \langle \hat{x}, A^* \hat{y} \rangle - \langle u, \hat{y} \rangle\}$$

$$\iff \begin{cases} y \in D(A^*) \text{ and } \exists x \in D(A) \text{ such that} \\ (*) \sup_{\hat{y}} \{H(x, \hat{y}) - \langle Ax, \hat{y} \rangle - \langle u, \hat{y} \rangle\} < \varepsilon + \inf_{\hat{x}} \{H(\hat{x}, y) - \langle \hat{x}, A^* y \rangle - \langle u, y \rangle\} \end{cases}$$

$$\iff y \in D(A^*) \text{ and } \exists x \in D(A) \text{ with } (**) (A^* y, -y) \in \partial_\varepsilon F(x, u + Ax).$$

If also (iii) Δ_0 is w.l.c. then, for each u such that

$$\bigcap_{x \in C \cap D(A)} \{y | \bar{H}(x, \cdot)^+(y) - \langle u + Ax, y \rangle > 0\} \text{ is a subspace,}$$

where $C := \{x | \bar{H}(x, \cdot) \text{ is somewhere finite}\}$ and $\bar{H}(x, \cdot)(y) := \text{usc } H(x, \cdot)(y)$, one has $p(u) \in \mathbb{R}$ and the "sup" defining $p(u)$ is actually "max."

Relation (*) (resp. (**)) for $\varepsilon = 0$ can be regarded as the abstract Hamiltonian (resp. Euler-Lagrange) system for the setting of Theorems V, IV (cf. [1]).

Main outlines of proof. (I) follows from (IV) by replacing A by $-A$ and putting $F(x,u) := h(x) + \psi_{\{0\}}(u)$, where generally $\psi_S := 0$ on S and $:= \infty$ off S . (V) is deduced from (IV) plus saddle function theory. (IV) splits into two cases: that of A continuous is proved by refining for the map $(x,u) \mapsto u$ the proof of (III) sketched below; that of A densely defined is deduced by applying

the first to $F_1(x,u) := F(x,u + Ax) + \psi_{D(A)}(x)$. Here, one proves as preliminary facts formulas for F_1^{0+} , F_1^* , $\partial_\epsilon F_1$. (II) involves two steps: the case $m = 2$ is proved by refining for the map $(x_1, x_2) \mapsto x_1 + x_2$ the proof of (III); then the continuous case of (IV) permits replacing f_2 by $f_2 \square \dots \square f_m$.

Sketch for (III). Observe $(Bk)^* \leq Ah$ always. Let $(Bk)^*(\bar{u}) < \infty$. Define a l.s.c. proper convex F by $F(w, w_1, w_2) := g_1(w - w_1) + g_2(w - w_2) - \langle w, \bar{z} \rangle$, for $W = W_1 = W_2 = V \times Y$, $Z = Z_1 = Z_2 = X \times U$, $\bar{z} = (0, \bar{u})$, $g_1 = \psi_{G(B)}$, $g_2 = k + \psi_Y$. One shows $\alpha := \inf_w F(w, 0, 0)$ and $\beta := \inf_{z_1, z_2} F^*(0, z_1, z_2)$ satisfy $\alpha = -\beta$, with \inf_w in β attained, if for $p(w_1, w_2) := \inf_w F(w, w_1, w_2)$ one has (a) $\alpha > -\infty$, (b) $(0, 0)$ and $\text{dom } p$ cannot be properly separated, and (c) \exists l.s.c. proper convex $\rho \geq p$ such that $\{(z_1, z_2) | \rho^*(z_1, z_2) < \gamma\}$, $\gamma \in \mathbb{R}$, are w.l.c. Now $\alpha = -(Bk)^*(\bar{u}) \Rightarrow$ (a), and (ii) \Leftrightarrow (b). For (c), take $\rho(w_1, w_2) := F(\bar{w} + w_1, w_1, w_2)$ for fixed $\bar{w} \in \text{dom } g_1$. Then $\rho(w_1, w_2) = g_2(\bar{w} + w_1 - w_2) - \langle w_1, \bar{z} \rangle + c$, $c := g_1(\bar{w}) - \langle \bar{w}, \bar{z} \rangle$, yields $\rho^*(z_1, z_2) = g_2^*(z) - \langle \bar{w}, z \rangle - c$ if $(\bar{z}, 0) + (z_1, z_2) = (z, -z)$ for some $z \in Z$, and $\rho^* = \infty$ otherwise. Then (i) \Rightarrow (c). Since $F^*(z, z_1, z_2) = h(-x_2)$ if $z = (x, u)$ and $z_1 = (x_1, u_1)$ satisfy $z + z_1 + z_2 = \bar{z}$, $(x_1, -u_1) \in G(A)$, $u_2 = 0$, and $F^* = \infty$ otherwise, $\beta = (Ah)(\bar{u})$. Thus, (1) holds. The ∂_ϵ assertions are proved with the aid of (1). For (2), observe $\text{epi } Ah$ lies between $(A \times I)\text{epi } h$ and its closure, where I is the identity on \mathbb{R} ; similarly for $\text{epi } A(h^{0+})$. Using (i), (ii) and T , $G(T) := G(A \times I) \cap (\text{epi } h \times U \times \mathbb{R})$, one proves $R(T) = (A \times I)\text{epi } h$ closed. Then, using (i), (ii) and S , $G(S) := G(I \times A \times I) \cap (\text{cl } H \times \mathbb{R} \times U \times \mathbb{R})$, where $H := \{a(1, x, \xi) | a > 0, (x, \xi) \in \text{epi } h\}$, one proves $R(S) = \{(\sigma, u, \mu) | \sigma > 0, (u, \mu) \in (A \times I)(\sigma \text{epi } h)\} \cup \{(0, u, \mu) | (u, \mu) \in (A \times I)^{0+}\text{epi } h\}$ is closed. Since $J := \{\sigma(1, u, \mu) | \sigma > 0, (u, \mu) \in (A \times I)\text{epi } h\}$ satisfies $J \subset R(S) \subset \text{cl } J = J \cup \{(0, u, \mu) | (u, \mu) \in 0^+(A \times I)\text{epi } h\}$, $R(S) = \text{cl } J$ follows, whence

$(A \times I)0^+ \text{epi } h = 0^+(A \times I) \text{epi } h$. The nontrivial half of this plus (ii) imply

$(A \times I) \text{epi } h$ contains no "vertical" line, completing (2).

Details and related results will appear elsewhere.

REFERENCES

- [1] J.-P. Aubin, Mathematical methods of game and economic theory, North-Holland, 1979.
- [2] V. Barbu and T. Precupanu, Convexity and optimization in Banach spaces, Sijthoff and Noordhoff, 1978.
- [3] I. Ekeland and R. Temam, Analyse convexe et problèmes variationnels, Dunod, 1974.
- [4] A. D. Ioffe and V. M. Tikhomirov, Teoriia ekstremal'nykh zadach, Nauka, 1974.
- [5] J. L. Joly and P.-J. Laurent, Stability and duality in convex minimization problems, Rev. Francaise Informat. Recherche Opérationnelle R-2, 1971, 3-42.
- [6] P.-J. Laurent, Approximation et optimisation, Hermann, 1972.
- [7] J. J. Moreau, Fonctionnelles convexes, lecture notes, Séminaire Équations aux dérivées partielles, Collège de France, 1966-67.
- [8] R. T. Rockafellar, Convex analysis, Princeton Univ. Press, 1970.
- [9] R. T. Rockafellar, Saddle-points and convex analysis, Differential Games and Related Topics (H. W. Kuhn and G. P. Szegö, eds.), North-Holland, 1971, 109-127.
- [10] R. T. Rockafellar, Conjugate duality and optimization, SIAM, 1974.

LM:scr

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2708	2. GOVT ACCESSION NO. AD-A144 716	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) QUASISTABLE PARAMETRIC OPTIMIZATION WITHOUT COMPACT LEVEL SETS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) L. McLinden		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041 MCS-8102684
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 5 - Optimiza- tion and Large Scale Systems
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.		12. REPORT DATE June 1984
		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASSIFICATION (of this report) UNCLIFIED
		15a. DECLASSIFICATION SCHEDULE SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) convex optimization weak local compactness perturbational duality noncoercive functionals recession theory abstract Hamiltonian system unbounded operators dual operations abstract Euler-Lagrange convex processes ϵ -subdifferentials system ϵ -solutions saddlepoint problems		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The well-known perturbational duality theory for convex optimization is refined to handle directly, in locally convex Hausdorff spaces, problems involving noncoercive convex functionals together with unbounded densely defined linear operators or, more generally, convex processes. The theory presented includes conjugacy, recession, and ϵ -subdifferential formulas for the two fundamental pairs of dual operations and also includes systematic treatment of ϵ -solutions.		

ATE
LMED
-8